
AD-A110 076

NAVAL AIR DEVELOPMENT CENTER WARMINSTER PA AIRCRAFT --ETC F/6 6/1
STUDIES ON THE MCCHANISM OF ACTION OF THE 'IN VITRO' POB SUB X --ETC(U)

OCT 61 M W SHMUKLER, M 6 ZAWRYT

UNCLASSIFIED NACC-81249-60 NL END $1 \sim 1$



REPORT NO. NADC-81244-60



70



STUDIES ON THE MECHANISM OF ACTION OF THE IN VITRO PGBx EFFECT

IV. THE EFFECT OF ORDER OF ADDITION OF PGB_X TO ASSAY SYSTEM

Herman W. Shmukler, Ph.D. Naval Air Development Center Aircraft and Crew Systems Technology Directorate Biochemistry Research Team Warminster, Pennsylvania 18974

> Margaret G. Zawryt, B.S. Hahnemann Medical College Philadelphia, Pennsylvania 19102

> > 16 October 1981

Phase Report Airtask No. F58527803 Work Unit No. EH810



APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED

Prepared for Office of Naval Research Department of the Navy Arlington, Virginia 22217

UTE FILE COPY

AD A110076

393532 01 26 82 022

NOTICES

REPORT NUMBERING SYSTEM - The numbering of technical project reports issued by the Navel Air Development Center is arranged for specific identification purposes. Each number consists of the Center acrenym, the calendar year in which the number was assigned, the sequence number of the report within the specific calendar year, and the official 2-digit correspondence code of the Command Office or the Functional Directorate responsible for the report. For example: Report No. NADC-78815-29 indicates the fifteeth Center report for the year 1978, and prepared by the Systems Directorate. The numerical codes are as follows:

CODE	OFFICE OR DIRECTORATE
00	Commander, Naval Air Development Center
01	Tachnical Director, Navel Air Development Center
02	Comptroller
18	Directorate Command Projects
20	Systems Directorate
30	Sensors & Avionics Technology Directorate
40	Communication & Navigation Technology Directorate
50	Softwere Competer Directorate
60	Aircraft & Crow Systems Technology Directorate
70	Planning Assessment Resources
80	Engineering Support Group

PRODUCT ENDORSEMENT -- The discussion or instructions concerning commercial products herein do not constitute an endorsement by the Government nor do they convey or imply the license or right to use such products.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

REPORT DOCUMENTATION PAGE	Read instructions Before completing f <u>orm</u>					
	3. RECIPIENT'S CATALOG NUMBER					
NADC-81244-60 AD-A110076						
4. TITLE (and Subtitie)	5. TYPE OF REPORT & PERIOD COVERED					
Studies on the Mechanism of Action						
of the in vitro PGBx Effect	Phase Report					
IV. The Effect of Order of Addition of PGBx	6. PERFORMING ORG. REPORT NUMBER					
to Assay System	8. CONTRACT OR GRANT NUMBER(s)					
Herman W. Shmukler and Margaret G. Zawryt	N00014-81-WR10070					
9. PERFORMING ORGANIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS					
Naval Air Development Center	1					
Aircraft & Crew Systems Technology Directorate	Airtask: @F58527803					
Warminster, PA 18974	Work Unit No.: EH810					
11. CONTROLLING OFFICE NAME AND ADDRESS	12. REPORT DATE					
Office of Naval Research	16 October 1981					
Department of the Navy Arlington, VA 22217	7					
14. MONITORING AGENCY NAME & ADDRESS(II different from Controlling Office)	18. SECURITY CLASS. (of this report)					
	UNCLASSIFIED					
	154. DECLASSIFICATION/DOWNGRADING					
16. DISTRIBUTION STATEMENT (of this Report)	<u> </u>					
is distribution statement for mis superly	12 3 1 1 N					
	•					
APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED						
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different fre	en Report)					
18. SUPPLEMENTARY NOTES						
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)						
Prostaglandin						
Oxidative Phosphorylation	100					
Rat Liver Mitochondria (RLM) PGB mib X	deg					
, GD						
20. ABSTRACT (Continue on reverse side if necessary and identify by block missher)						
The mechanism of action of PGBx in RLM oxidative phosphorylation was studied by						
varying the sequence of addition of PGBz When PGBz was added to RIM pre-						
exposed to hypotonic media at 270 no restoration of phosphorylation was observed.						
When RLM were exposed to hypotonic conditions for 30 seconds before the addition						
of PGB _X and subsequently exposed to hypotonic conditions, the resulting phos-						
phorylation was 60% of that observed in the normal system. From these results it is concluded that PGB_{x} functions in the in vitro PGB_{x} RLM assay system as a						
It is concluded that PGB _N functions in the in vitro "protective" agent rather than a "restorative" agen	Cropy Jers sessa sastem es e					
	<u> </u>					
DD 1 JAN 73 1473 EDITION OF 1 NOV 68 IS OSSOLETE	UNCLASSIFIED					

ECURITY CLASSIFICA	HON OF THIS PAGE	(with bits and				
			•	•		
					-	
					-	
			•			
				,		

SECURITY CLASSIFICATION OF THIS PASE(When Dots Enters

TABLE OF CONTENTS

	Page
INTRODUCTION	2
EXPERIMENTAL	2
DISCUSSION	3
TABLE I	4
FIGURE 1	5
REFERENCES	6

COPY INSPECTED

A

INTRODUCTION

The possible use of PGB_X (1, 2, 3) as a therapeutic agent in the treatment of human ischemic diseases was suggested by the favorable results from only a few animal experiments (4, 5, 6, 7, 8, 9, 10). Obviously before any human trials may be undertaken much more animal testing must be completed, and in addition, certain basic information concerning the mechanism of action of this drug must be known. At this stage of our knowledge of PGB_X biochemistry, the possibility of successful elucidation of the in vivo mechanism of action appears remote. However, since the elucidation of the in vitro mechanism of PGB_X might reveal an insight as to the in vivo mechanism, studies have been underway in this laboratory to define the in vitro PGB_X effect.

Recently we reported that PGB_X ineracts with BSA (11) to form a complex that was inactive in the <u>in vitro</u> PGB_X assay system (2, 3). Some of the experimental evidence published in that report was based on altering the sequence of the addition of PGB_X and BSA in the assay system so that the PGB_X and BSA could react before the addition of RIM. The results of these experiments then suggested the importance of the sequence of addition of the reactants of the test system, and that by studying this in detail some information concerning the mechanism of the <u>in vitro</u> PGB_X effect might be realized.

EXPERIMENTAL

<u>Materials and Methods</u>: PGB_X Type II was synthesized and assayed for the <u>in vitro</u> PGB_X effect as reported by Polis et al (2, 3). RLM was isolated and stored at 0^0 as described previously (2, 3). In this study the RLM were aged three days before use. All analytical methods used in this study were described previously (2, 3).

Note: Abbreviations used in this report are: RLM, rat liver mitochondria; BSA, bovine serum albumin; Pi, inorganic phosphate

Results: The effect of varying the order of addition of PGBx to the in vitro PGBx assay system was studied by measuring the changes in phosphorylating ability of RLM under various experimental conditions. Table I lists the composition and experimental conditions for the in vitro PGBx assay system (2). In the normal PGBx assay system, RLM were added to the hypotonic medium (Mixture A, Table I) containing the requisite amount of PGB, and the mixture then incubated for the required time period. Phosphate acceptor (Mixture B, Table I) was then added and the mixture again incubated for 20 minutes. At the end of this time period the solution was deproteinized and the Pi esterfied was measured. The extent of phosphorylation by this system was compared to that resulting from the following changes in experimental protocol: (a) RLM were incubated in Mixture A in the absence of $PGB_{\mathbf{x}}$ for the required time period; at the end of this incubation period PGBx was added followed by the addition of Mixture B; the assay was then continued as above. (b) RLM was added to Mixture A and incubated for 30 seconds; PGBx was then added and the hypotonic incubation continued for the required time period; at the end of the incubation period Mixture B was added and the assay continued as above. Figure I shows the results of these tests. The curves shown are described in the legend of the figure. The normal assay curve of $PGB_{\mathbb{X}}$ concentration shows the usual biphasic response with the maximum phosphorylation in the range of 2-10 $\mu g/ml$ of reaction. When RLM were incubated in Mixture A in the absence of PGB_X , the subsequent addition of PGB_X did not restore the phosphorylation ability of the degraded RLM. When PGB_{X} was added to the RLM incubated in Mixture A for 30 seconds, a biphasic phosphorylation curve was observed, however the maximum phosphorylation ability was only about 2/3 of the normal assay and in addition was shifted so that 10-20 µg PGBx/ml reaction were required.

DISCUSSION

In earlier reports on the physiological mechanism of PGB_X action in both in vitro and in vivo systems it was suggested that PGB_X functioned to reactivate damaged mitochondria (1, 2, 3, 8, 12) and thus reverse the pathological effects of ischemia. This interpretation appears contrary to the results reported in this study with isolated RLM subjected to hypotonic degradation. The results reported here show that RLM exposed to hypotonic media at 27° show a reduced capacity for oxidative phosphorylation. However when RLM are exposed to hypotonic media at 27° containing the required amount of PGB_X , they maintain their capacity for oxidative phosphorylation. This phenomenon is the in vitro PGB_X effect on RLM. In contrast when RLM are exposed to hypotonic media at 27° , the subsequent addition of PGB_X has no effect on the recovery of phosphorylation activity of these RLM.

From the above results it may be concluded that PGB_X functions in the in vitro assay system as a protective agent for RLM capacity for phosphory-lation, rather than a "restorative" agent that reverses the degradative effects of exposure of RLM to hypotonic conditions.

TABLE I

The Composition of the Medium for the Demonstration of the $\text{PGB}_{\mathbb{X}}$. Effect on Mitochondrial Oxidative Phosphorylation


Order of Addition	Mitochondrial Degrading Medium	Reaction Mixture
Water	1.55 ml	1.55 ml
Phosphate Buffer pH 7.35	4.98 mM	4.55 mM
α-Ketoglutarate pH 7.35	14.93 mM	13.64 mM
MgSO4	4.98 mM	4.55 ™
Aged Mitochondria	1.99 mg/ml	1.82 mg/ml
Sucrose*	5.97 mM	5.45 mM
EDTA*	0.010 mM	0.009 mM
AMP		2.27 mM
ADP		2.27 mM
KC1		45.45 mM
Bovine Serum Albumin	****	0.68 mg/ml

Total Volume: 2.20 ml

Temperature: 280

Degradation Time: 5-20 minutes Reaction Time: 20 minutes

*Added with mitochondris



REFERENCES

- Polis, B. D., A. M. Grandizio and E. Polis: Some in vitro and in vivo
 Effects of a New Prostaglandin Derivative. Neurohumoral and Metabolic
 Aspects of Injury. Adv. in Exper. Med. and Biol., Plenum Press, New York
 33:213-220 (1973).
- 2. Polis, B. D., S. F. Kwong, E. Polis and G. Nelson: Studies on PGB_X : A Polymeric Derivative of Prostaglandin B₁: I Synthesis and Purification of PGB_X . Report No. NADC-78235-60 (1978).
- Polis, B. D., E. Polis and S. F. Kwong: Protection and Reactivation of Oxidative-Phosphorylation in Mitochondria by a Stable Free-Radical Prostaglandin Polymer (PGB_x). <u>Proc. Natl. Acd. Sci.</u>, 76:1598-1602 (1979).
- Angelakos, W. T., B. D. Polis and R. L. Riley: Recovery After Coronary Ligation and Fibrillation in Primates Treated with a Prostaglandin Derivative. 47th Scientific Sessions American Heart Association, Abstracts, Dallas, Texas (1974).
- Angelakos, E. T., B. D. Polis and R. L. Riley: Protection by a Prostaglandin Derivative from Mortality after Coronary Ligation from Ventricular Fibrillation in Primates. 6th International Congress of Pharmacology Abstracts, Helsinki, Finland (1975).
- Angelakos, E. T., R. L. Riley and B. D. Polis: Recovery of Monkeys from Cardiogenic Shock After Myocardial Infarction with Ventricular Fibrillation. Effects of PGB_x. Report No. NADC-77308-60 (1977).
- Kolata, R. J.: The Effect of PGB_X on Neurological Recovery from Cerebral Ischemia in Rabbits. <u>Masters Thesis</u>, <u>Univ. of Penna. Veterinary School</u> (1977).
- 8. Kolata, R. J. and B. D. Polis: Facilitation of Recovery from Ischemic Brain Damage in Rabbits by Polymeric Prostaglandin PGB_x, a Mitochondrial Protective Agent. Physiol. Chem. and Physics 12, 551 (1980).
- Yamazaki, H., M. M. Bodenheimer, V. S. Banka, J. Lewandowski and R. H. Helfant: The Effect of a New Prostaglandin (PGB_x) on Length-tension Relationship Following Partial Coronary Occlusion and Reperfusion.
 Amer. Heart Assn., Abstracts 51st Scientific Session, Dallas, Texas, (1978).

- 10. Moss, G., T. Magliochetti and R. Quarmby: Immediate Restoration of Central Nervous System Autonomic Cardio-pulmonary Control: Survival of "Lethal" Cerebral Hypoxia by Treatment with $PGB_{\mathbf{X}}$. Surgical Forum, 21:513 (1978).
- 11. Shmukler, H. W., M. G. Zawryt, W. Feely and E. Polis: The Interaction of PGB_X with Bovine Serum Albumin. Physiol. Chem. and Physics, 13, 241 (1981).
- 12. Polis, B. D. and E. Polis: Novel Prostaglandin Derivatives, certain in vivo and in vitro Effects thereof and Processes for the Preparation of Same.
 United States Patent (19) (11) 4, 153, 808, 8 May 1979.

DATE FILMED

DTIC